


Claims

1. A compound of formula (I):






5



(I)

wherein:

R¹ represents a group selected from:



10 each ring of which optionally contains a further heteroatom N,
Z represents an optional substituent halogen,
alk represents alkylene or alkenylene,
T represents S, O or NH;

15 R² represents hydrogen, -C₁₋₆alkyl, -C₁₋₃alkylCONR^aR^b, -C₁₋₃alkylCO₂C₁₋₄alkyl, -CO₂C₁₋₄alkyl or -C₁₋₃alkylCO₂H;

R^a and R^b independently represent hydrogen, $-C_{1-6}$ alkyl, or together with the N atom to which they are bonded form a 5-, 6- or 7- membered non-aromatic heterocyclic ring optionally containing an additional heteroatom selected from O, N or S, optionally substituted by $-C_{1-4}$ alkyl, and optionally the S heteroatom is substituted by O, i.e.
5 represents $S(O)_n$;

n represents 0-2;

X represents phenyl or a 5- or 6- membered aromatic heterocyclic group containing at
10 least one heteroatom selected from O, N or S, each of which is optionally substituted by 0-2 groups selected from: halogen, $-C_{1-4}$ alkyl, $-C_{2-4}$ alkenyl, $-CN$, $-CF_3$, $-NR^aR^b$, $-C_{0-4}alkylOR^e$, $-C(O)R^f$ and $-C(O)NR^aR^b$;

R^e represents hydrogen or $-C_{1-6}$ alkyl;


15 R^f represents $-C_{1-6}$ alkyl;

Y represents phenyl or a 5- or 6- membered aromatic heterocyclic group containing at least one heteroatom selected from O, N or S, each of which is substituted by a group $-C_{1-2}alkylNR^cR^d$.
20

R^c and R^d , together with the nitrogen atom to which they are bonded, form a 4-membered heterocyclic ring optionally substituted by halogen, OH or $-OC_{1-6}alkyl$, or a 5- or 6-membered non-aromatic heterocyclic ring substituted by OH, $-OC_{1-6}alkyl$ or 1 to 2
25 halogens, with the proviso that the substituent is not attached to a ring carbon atom adjacent to a heteroatom;

and/or pharmaceutically acceptable derivative thereof.

30 2. A compound according to claim 1 wherein R^1 represents a group selected from:



each ring of which optionally contains a further heteroatom N,
 Z represents an optional substituent halogen,
 alk represents alkylene or alkenylene,
 T represents S, O or NH.

5 and/or pharmaceutically acceptable derivative thereof.

3. A compound according to claim 1 or claim 2 wherein R² represents hydrogen and/or pharmaceutically acceptable derivative thereof.

10 4. A compound according to any one of claims 1-3 wherein X represents phenyl or a 5 or 6 membered aromatic heterocyclic group containing at least one heteroatom selected from O, N or S, each of which is optionally substituted by 0-2 groups selected from: halogen, -C₁₋₄alkyl or -NR^aR^b and/or pharmaceutically acceptable derivative thereof.

15 5. A compound according to any one of claims 1-4 wherein Y represents a 5 or 6 membered aromatic heterocyclic group containing at least one heteroatom selected from O, N or S, each of which is substituted by a group -CH₂NR^cR^d and/or pharmaceutically acceptable derivative thereof.

20 6. A compound according to claim 1 selected from:

(1E)-N-(1-{4-[2-(1-Azetidinylmethyl)-1H-imidazol-1-yl]-2-fluorophenyl}-2-oxo-3-pyrrolidinyl)-2-(5-chloro-2-thienyl)-1-propene-1-sulfonamide;

N-(1-{4-[2-(1-Azetidinylmethyl)-1H-imidazol-1-yl]-2-fluorophenyl}-2-oxo-3-pyrrolidinyl)-2-(5-chloro-2-thienyl)ethanesulfonamide;


25 N-((3S)-1-{4-[2-(1-Azetidinylmethyl)-1H-imidazol-1-yl]-2-fluorophenyl}-2-oxo-3-pyrrolidinyl)-6-chloro-1-benzothiophene-2-sulfonamide.

(E)-2-(5-Chloro-2-thienyl)-N-[1-(2-fluoro-4-{2-[(3-fluoro-1-pyrrolidinyl)methyl]-1H-imidazol-1-yl}phenyl)-2-oxo-3-pyrrolidinyl]ethanesulfonamide;


(1E)-2-(5-Chloro-2-thienyl)-N-[1-(2-fluoro-4-{2-[(3-fluoro-1-pyrrolidinyl)methyl]-1H-imidazol-1-yl}phenyl)-2-oxo-3-pyrrolidinyl]-1-propene-1-sulfonamide;

30

- 6-Chloro-*N*-[1-(2-fluoro-4-{2-[(3-fluoro-1-pyrrolidinyl)methyl]-1*H*-imidazol-1-yl}phenyl)-2-oxo-3-pyrrolidinyl]-1-benzothiophene-2-sulfonamide;
 6-Chloro-*N*-[1-[2-fluoro-4-(2-{[3-(methyloxy)-1-azetidiny]methyl})-1*H*-imidazol-1-yl)phenyl]-2-oxo-3-pyrrolidinyl]-1-benzothiophene-2-sulfonamide formate;
 5 and/or pharmaceutically acceptable derivative thereof.
7. A compound according to any of claims 1-6 and/or pharmaceutically acceptable derivative thereof for use in therapy.
- 10 8. A pharmaceutical composition comprising a compound according to any of claims 1-6 and/or pharmaceutically acceptable derivative thereof together with at least one pharmaceutical carrier and/or excipient.
9. Use of a compound according to any of claims 1-6 and/or pharmaceutically acceptable
 15 derivative thereof for the manufacture of a medicament for the treatment of a patient suffering from a condition susceptible to amelioration by a Factor Xa inhibitor.
10. A method of treating a patient suffering from a condition susceptible to amelioration by a Factor Xa inhibitor comprising administering a therapeutically effective amount of a
 20 compound according to any of claims 1-6 and/or pharmaceutically acceptable derivative thereof.
11. A process for preparing a compound of formula (I) which comprises:
- 25 (a) reacting a compound of formula (II) or an acid addition salt thereof with a compound of formula (III) where V is a suitable leaving group:



(II)



(III)

(b) by reacting compounds of formula (I) where R^2 is hydrogen with compounds of formula (XI):

5



wherein R^2 is $-C_{1-6}alkyl$, $-C_{1-3}alkylCONR^aR^b$, $-C_{1-3}alkylCO_2C_{1-4}alkyl$, or $-CO_2C_{1-4}alkyl$ and T is a suitable leaving group, optionally followed by removal of the alkyl protecting group
10 where appropriate.